

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Marvasi, Massimiliano, Casillas-Santiago, Lianne M., Henríquez, Tania and Casillas-Martinez, Lilliam (2017) Involvement of *etfA* gene during CaCO₃ precipitation in *Bacillus subtilis* biofilm. *Geomicrobiology Journal*, 34 (8) . pp. 722-728. ISSN 0149-0451 [Article]
(doi:10.1080/01490451.2016.1248254)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/23145/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

1 **Involvement of *etfA* gene during CaCO₃ precipitation in *Bacillus subtilis* biofilm**

2
3 *Massimiliano Marvasi*¹, *Lianne M. Casillas-Santiago*², *Tania Henríquez*^{1,3} and *Lilliam Casillas-*
4 *Martínez*^{2*}

5
6 ¹ Middlesex University London. The Burroughs, London NW4 4BT, UK.

7 ² University of Puerto Rico in Humacao, Biology Department, CUH 100 Carr. 908, Humacao,
8 00791, PR (USA). Email: lcasillasm@gmail.com. *Corresponding author.

9 ³ University of Chile. Department of Microbiology and Mycology, University of Chile, Institute
10 of Biomedical Sciences (ICBM).

11 **Key words:** *etfA*, *Bacillus subtilis*, biofilm, calcium carbonate, FAME analysis, dipicolinic acid,
12 teichoic acids

Abstract

The *eftA* gene in *Bacillus subtilis* has been suggested to be involved in the oxidation/reduction reactions during fatty acid metabolism. Interestingly *eftA* deletion in *B. subtilis* results in impairment in CaCO_3 precipitation on the biofilm. Comparisons between the wild type *B. subtilis* 168 and its *eftA* mutant during *in vitro* CaCO_3 crystal precipitation (calcite) revealed changes in phospholipids membrane composition with accumulation of up to 10% of *anteiso*- $\text{C}_{17:0}$ and 11% *iso*- $\text{C}_{17:0}$ long fatty acids. Ca^{2+} nucleation sites such as dipicolinic acid and teichoic acids seem to contribute to the CaCO_3 precipitation. *eftA* mutant strain showed up to 40% less dipicolinic acid accumulation compared with *B. subtilis* 168, while a *B. subtilis* mutant impaired in teichoic acids synthesis was unable to precipitate CaCO_3 . In addition, *B. subtilis eftA* mutant exhibited acidity production leading to atypical flagella formation and inducing extensive lateral growth on the biofilm when grown on 1.4% agar. From the ecological point of view, this study shows a number of physiological aspects that are involved in CaCO_3 organomineralization on biofilms.

Introduction

Electron transfer flavoproteins (ETFs) are *alpha beta*-heterodimers involved in the electron transfer during the oxidation/reduction reactions that takes place in fatty acid metabolism of eukaryotic mitochondria as well as bacteria (Tsai and Saier, 1995). Comparison of the amino acid sequences among all available ETFs and ETF-like proteins revealed the existence of different groups of ETFs. The groups are related to acyl-CoA dehydrogenases or enclose ETF-like proteins that are involved in growth under anaerobic condition (Weidenhaupt et al. 1996). In obligate anaerobic bacteria such as *Clostridium acetobutylicum*, the co-expression of *etfA* and *etfB* was essential for the butyryl-CoA dehydrogenase (BCD) activity of fatty acids biosynthesis (Boynton et al. 2006; Inui et al. 2008). EtfA-B proteins of *Bacillus subtilis* show 64% and 57% similarity with EtfA and EtfB proteins from *C. acetobutylicum*, respectively. However in *B. subtilis* 168 *etfA-B* functions are only putative, the genes are not essential for growth and do not results in differential cell morphology (Kobayashi et al. 2003; Barabesi et al. 2007). In *B. subtilis*, *etfA* is regulated by the central regulator *fadR*. The *fadR* regulon is involved in the β -oxidation cycle comprises five operons: *lcfA-fadR-fadB-etfB-etfA*, *lcfB*, *fadN-fadA-fadE*, *fadH-fadG*, and *fadF-acdA-rpoE* (Matsuoka 2007; Tojo et al 2011).

Interesting, when *B. subtilis* 168 is grown on B4 precipitation medium supplemented with calcium acetate (0.25% w/v) it produces CaCO_3 calcite crystals on biofilm after 6-7 day of incubation at 39°C (Barabesi et al. 2007). Barabesi and collaborators (2007) generated a mutant strain in *etfA* gene which cannot precipitate CaCO_3 crystals on its biofilm. In summary, *etfA* mutation is linked to fatty acid metabolism but its overall effects in cell physiology during CaCO_3 precipitation are still unknown. The main goal of this study is the elucidation of the physiological responses altered by the *etfA* mutation in *B. subtilis* during CaCO_3 precipitation in

vitro.

Materials and Methods

Bacterial strains and growth conditions. Strains used in this study were *Bacillus subtilis* 168 (Anagnostopoulos and Spizizen, 1961), *B. subtilis* 168 mutated in *etfA* gene (Barabesi et al. 2007), and *B. subtilis* EB1451 [*hisA1 argC4 metC3 tagO::Erm^r*] impaired in teichoic acid synthesis. *B. subtilis* cultures were routinely grown on Nutrient Agar (Oxoid) and standard B4 precipitation medium (0.4% yeast extract, 0.5% dextrose, 0.25% calcium acetate and 1.4% agar when solid medium was prepared) (Boquet et al. 1973). For studies related to flagella formation, modified B4 medium lacking calcium acetate was used to avoid calcium interferences during chemotaxis (Ordal et al. 1983; Herbaud et al. 1998). Modified buffered B4 medium was prepared by addition of 1.2% TRIS-HCl stabilized at pH 7.3 with 2N HCl.

Unless specified otherwise, biofilms were grown on plates incubated at 39° C inside a plastic bag to prevent dehydration.

Membrane Fatty Acid Methyl Esters (FAME) analysis (Eder, 1995). Cultures from *B. subtilis* 168 strain and *etfA* mutant were normalized to OD₆₀₀=0.5 and 200 µl were inoculated on B4 plates. Plates were incubated 8 hours, 7 and 30 days at 37°C and biofilms were scraped and subjected to whole cell fatty acid methyl esters analysis. All the experiments were conducted in triplicates. FAME analysis samples were initially homogenized and subjected to saponification at 100°C with 1 ml of methanolic NaOH (15% NaOH in 50% methanol) followed by esterification of the fatty acids at 80°C with 2 ml of 3.25 N HCl in 46% (vol/vol) methanol. FAMES were extracted into 1.25 ml of 1:1 (vol/vol) methyl-tert-butyl etherhexane followed by

an aqueous washing of the organic extract with 3 ml of 1.2% (wt/vol) NaOH methylation and extraction as described by Kidd Haack et al. (1994). Profiles for total fatty acids methyl esters (FAME) in each sample were generated after gas chromatography using a 25 m x 0.2 mm phenyl methyl silicone fused silica capillary column. The chromatography was conducted using an HP 5890 series II gas chromatograph (Hewlett Packard, Palo Alto, California) with the oven temperature ramping up from 170 °C to 270 °C at 5 °C per minute and a final step of 300 °C for 2 min. Individual compounds were quantified by flame ionization detector (FID) in response to internal standards added prior to CG analyses. Fatty acids between 9 and 20 carbons in length were analyzed and identified using the Sherlock MIS software (MIDI, Inc., Newark, Delaware). Final (%) of total fatty acids resulted from an automatic comparison of the composition of the unknown strain to a stored database using a covariance matrix, principal component analysis and pattern recognition software. The covariance matrix takes into account the mole-for-mole relationship of the conversion of one fatty acid to another (e.g. 16:0 to 16:1 due to action of a desaturase), which might occur in relation to a temperature shift or age difference. The pattern recognition software uses calculations of cross terms (e.g. ratios between fatty acid amounts) in addition to the principal component base. Experiments were conducted in triplicate.

Dipicolinic acid (DPA) assays. *B. subtilis* biofilms were grown in B4 solid medium and scraped from the plates after 12 hours, 1, 2, 3, 4 days of incubation at 39 °C until 30 mg of wet weight was achieved. Due to the formation of CaCO₃ crystals on *B. subtilis* 168, biofilms samples were centrifuged at 10,000xg for 10 min and the upper phases were collected into a new pre-weighted tube. These samples were resuspended in 1 ml of distilled water and measurements of total DPA were performed according to Nicholson and Setlow, 1990. Experiments were conducted in triplicate.

Microscopic analysis. To monitor crystal formation in the biofilms we used a stereo microscope Leica ES2. An Optical Nikon Eclipse E400 microscope was used for microscopic analysis of flagella content. Samples from the *B. subtilis* biofilms were stained using the BD Flagella Stain Dropper kit according to the manufacture's manual (Becton, Dickinson and Company). To record images microscopes were integrated into a Nikon Spot Insight digital camera. Experiments were conducted from three different sections of the biofilm and at least three different plates were tested.

Results

B. subtilis eftA mutant accumulates up to 21% more long fatty acids (C_{17:0}) when compared to B. subtilis 168 during in vitro CaCO₃ precipitation

B. subtilis 168, as well as other soil bacteria, is capable of CaCO₃ (calcite) formation if grown in B4 precipitating medium starting from day 7th at 39°C (Figure 1, A). On the opposite, mutation in the *eftA* gene prevented calcite crystal formation in *eftA B. subtilis* mutants grow under similar conditions. (Figure 1, B) (Barabesi et al. 2007). To measure the variation of total lipids within the plasma membrane during CaCO₃ precipitation, *B. subtilis* 168 and *eftA* mutant were analysed via Fatty Acid Methyl Ester (FAME) analysis (Table I). During the first 8 hours of growth on B4 medium no differences were reported among the *B. subtilis* 168 and *etfA* mutant. However, after one week and one month of incubation, differences were reported for C_{15:0} and C_{17:0} (Table I, Figure 2). *B. subtilis* 168 accumulated up to 13 % more of *anteiso*-C_{15:0} when compared with *etfA* mutant. On the contrary, longer fatty acids *anteiso*-C_{17:0} and *iso*-C_{17:0} accumulated in *etfA* mutant, up to 10% and 11%, respectively, after one month of incubation (Figure 2).

128
129 *Ca²⁺ chelants such as dipicolinic acid and teichoic acid contribute to CaCO₃ precipitation in B.*
130 *subtilis.*

131 In *Clostridium perfringens*, EftA catalyses the formation of pyridine-2,6-dicarboxylic acid
132 (commonly named dipicolinic acid or DPA) (Osburn et al. 2010). Interestingly, dipicolinic acid
133 forms a complex with calcium ions within the endospore core (Hintze and Nicholson 2010).
134 Consequently, we compared the levels of DPA of *B. subtilis* 168 and the *B. subtilis eftA* mutant
135 biofilms during growth on B4 medium. Although initially DPA accumulation is similar for both
136 strains, after 42 hours dipicolinic acid concentration was 40% lower in the *B. subtilis eftA* mutant
137 when compared with the wild type (Figure 3). Considering another Ca²⁺ chelator in addition to
138 dipicolinic acid, we also wondered whether the external charge of teichoic acids could have a
139 role in CaCO₃ precipitation. *B. subtilis* EB1451 strain mutated in *tagO* gene and unable to
140 produce any wall teichoic acid was tested (D'Elia et al. 2006). When *B. subtilis* EB1451 biofilm
141 was formed on B4 solid medium and incubated at 37°C for 21 days, CaCO₃ precipitation was
142 impaired.

143
144 *B. subtilis eftA mutant is capable of flagella formation when compared to B. subtilis 168*
145 Mutation of *eftA* is known to cause increasing of acidity of the *B. subtilis* biofilm due to an
146 excess of proton (H⁺) extrusion from the cell (Marvasi et al. 2010). Release of H⁺ across the
147 cytoplasmic membrane is coupled with rotation of most bacterial flagella including those from *B.*
148 *subtilis* (Matsuura et al. 1979; Ito et al. 2005). To analyse phenotypic changes in flagella
149 formation in *B. subtilis* caused by *eftA* mutation, calcium acetate was eliminated from B4
150 medium as calcium tunes *Bacillus* sp. chemotaxis (Ordal et al. 1983; Herbaud et al. 1998).

151 Interestingly, the biofilm morphology of *B. subtilis eftA* mutant showed extensive lateral growth
152 (Figure 4, panel A) indicative of the presence of highly motile cells. Lateral growth was not
153 observed for *B. subtilis* 168 strain (Figure 4, panel B). Further microscopic observations on the
154 bacteria growing in the edges of the lateral branches formed by *B. subtilis eftA* mutant revealed
155 the presence of flagella (Figure 3, panel A). To our knowledge, flagella formation at 1.4% of
156 agar has not been previously reported for any *Bacillus subtilis* strain (Senesi et al. 2004).
157 To determine if extensive lateral growth was associated with an excess of protons (H^+) extrusion
158 previously reported in the *B. subtilis eftA* mutant, strains were grown on buffered B4 medium to
159 sequester H^+ from the medium. Lateral growth and flagella formation were not observed when
160 the *B. subtilis* strains were grown on B4 pH 7.3 buffered medium (Figure 4, panel C, D).

163 Discussion

164 A number of bacteria are capable to form minerals by biologically induced mineralization (BIM)
165 processes in which the minerals generally nucleate and grow extracellularly as a result of
166 bacterial metabolic activities (Visscher et al. 2010). Calcium carbonate precipitation in *B. subtilis*
167 is one of these BIM processes that is poorly understood at the physiological and genetic levels.
168 Recently, Barabesi et al. 2007 isolated a mutant strain in *etfA* gene impaired in $CaCO_3$ crystals
169 formation on its biofilm. Further analysis showed that the impairment was originated due to an
170 excess of production of protons from the biofilm (Marvasi et al. 2010).
171 *etfA* participate in fatty acid β -oxidation cycle, affecting the utilization of palmitic acid as
172 representative of long chain fatty acids and the accumulation of branched and straight long chain
173 acyl-CoAs in *B. subtilis* (Matsuoka et al. 2007). The FAME profiles during $CaCO_3$ precipitation

of *B. subtilis* 168 and *etfA* mutant biofilms (Table I) indicated that mutations in *etfA* caused an accumulation of *anteiso*-C_{17:0} and *iso*-C_{17:0}, compared to the shortest *anteiso*-C_{15:0}, *iso*-C_{15:0} after the first week of incubation. The percentage of fatty acids accumulation in both *B. subtilis* 168 and *etfA* mutant does not change after 8 hours, while the difference are clearly visible after 1 week where *anteiso*-C_{15:0} decreases and *iso*-C_{17:0} and *anteiso*-C_{17:0} increase in *B. subtilis etfA* (Figure 2). Accumulation of long chain fatty acids may be explained due to *etfA* deletion, which inhibits YsiA, a central regulator in fatty acid degradation, causing the accumulation of branched and straight long chain acyl-CoAs (Matsuoka et al. 2007). The link between plasma membrane fatty acids composition and CaCO₃ precipitation may be ascribed due to changes of membrane fluidity which is one of the responses that the bacteria uses to cope with environmental stress (Beranová et al. 2008). We can speculate that such changes may lead to the change of number of calcium nucleation sites. Indeed, membrane fluidity has been linked with the control of important calcium nucleation sites such as DPA and teichoic acids (Cowan et al. 2004, Zhang and Rock 2008). As previously mentioned, DPA and teichoic acids are able to bind calcium. DPA is a main key player of spore resistance to many environmental stresses during long periods of dormancy and in *Clostridium perfringens*, EtfA protein catalyses the formation of DPA (Huang et al. 2007; Orsburn et al. 2010). DPA analysis showed that *B. subtilis etfA* biofilm exhibits a 40% less DPA when compared with the *B. subtilis* 168 (Figure 3). Dipicolinic acid forms a complex with calcium ions within the endospore core and mutants in dipicolinic acid fail in accumulate calcium on the cell surface (Hintze and Nicholson 2010, Hanson et al 1978). The reduction of DPA in the mutant could contribute to the impairment of CaCO₃ precipitation at neutral pH. Teichoic acids could also be responsible of mineral precipitation, serving as nucleation site for

calcium. To confirm this hypothesis, *B. subtilis* mutant EB1451 strain (D'Elia et al. 2006) unable to produce any wall teichoic acid was totally impaired in CaCO_3 precipitation. Current research has shown the role of teichoic acids as nucleation sites or Ca^{2+} -carriers. It is well known that the cell wall of Gram positive bacteria presents teichoic acids covalently bound to the cell wall or anchored in the cytoplasmic membrane as lipoteichoic acid (Thomas et al. 2014). The polyphosphate groups of teichoic acid provide one-half of the metal binding sites for calcium (Thomas et al. 2014), while the other Ca^{2+} bind the carboxyl units of peptidoglycan. According with Thomas and collaborators (2014) curvature of Scatchard plots showed two regions of binding affinity: Region I $K_A = (1.0 \pm 0.2) \times 10^6 \text{ M}^{-1}$ and Region II $K_A = (0.075 \pm 0.058) \times 10^6 \text{ M}^{-1}$. Binding capacity for both regions (η_2) is $0.70 \pm 0.04 \text{ } \mu\text{mol/mg}$ for Ca^{2+} . Calcium binding constants are highly dependent from the calcium concentration and cell wall type (Thomas et al. 2014). Experiments on crystal formation *in vitro* show controversial data: Adsorption of hydroxyapatite to [3H]-lipoteichoic acid effects positively crystal growth (Damen et al. 1994) while in oral streptococci, lipoteichoic acids inhibited Ca^{2+} -phosphate precipitation in a distinct pH-range just above the acid solubility of the mineral by complexing Ca^{2+} -ions (Bergmann et al 1991).

With reference to the flagella formation experiment, it was carried out mainly to confirm the acidic phenotype of *etfA*. Prior studies revealed that the *etfA* mutant extruded $0.7 \text{ moles H}^+\text{L}^{-1}$ more with respect to *B. subtilis* 168 strain on B4 medium and that the extrusion of protons was contributing to the impairment of CaCO_3 (Marvasi et al. 2010). Interestingly, *B. subtilis etfA* mutant exhibits flagella formation when grown on B4 media plates with agar concentration 1.4% with extensive lateral growth (Figure 4). However, if the *B. subtilis etfA* mutant was plated on B4 buffered medium pH 7.3 such lateral grow and flagella formation was arrested. Protons extrusion

across the cytoplasmic membrane is coupled with rotation of most bacterial flagella including *B. subtilis* (Matsuura et al. 1979; Ito et al. 2005). These data seem to suggest that in *B. subtilis etfA* mutant the proton excess could induce flagella formation through an indirect system. Once the protons are sequestered (such as it is the case in the buffered B4 medium), *B. subtilis* flagella formation is arrested. In *B. subtilis* strain PB1831 strain, Senesi and collaborators (2004) reported lack of flagella formation during growth at agar concentrations ranging from 0.5 to 2%. All these data, show how *etfA* mutation, which has been previously characterized as the main responsible for the impairment of CaCO₃ precipitation, leads to several different metabolic divergences when compared with the wild type that ultimately contribute in the prevention of CaCO₃ organomineralization in *B. subtilis* biofilm. DPA and teichoic acids could contribute to organomineralization in many other bacterial species (Marvasi et al. 2011).

Bibliography

- Anagnostopoulos C, Spizizen J. 1961. Requirements for transformation in *Bacillus Subtilis*. J Bacteriol 81(5):741-6.
- Barabesi C, Galizzi A, Mastromei G, Rossi M, Tamburini E, Perito B. 2007. *Bacillus subtilis* gene cluster involved in calcium carbonate biomineralization. J Bacteriol 189(1):228-35.
- Bergmann JE, Gülzow HJ, Lewandowski T. 1991. Oral streptococci affecting Ca-phosphate precipitation. Dtsch Zahnarztl Z 46(8):561-2.
- Boquet E, Boronat A, Ramos-Cormenzana A. 1973. Production of calcite (calcium carbonate) crystals by soil bacteria is a general phenomenon. Nature 246:527-529.
- Boynton ZL, Bennet GN, Rudolph FB. 1996. Cloning, sequencing, and expression of clustered

243 genes encoding beta-hydroxybutyryl-coenzyme A (CoA) dehydrogenase, crotonase, and
 244 butyryl-CoA dehydrogenase from *Clostridium acetobutylicum* ATCC 824. J Bacteriol
 245 178(11):3015-24.
 246 Cowan AE, Olivastro EM, Koppel DE, Loshon CA, Setlow B, Setlow P. 2004. Lipids in the
 247 inner membrane of dormant spores of *Bacillus* species are largely immobile. Proc Natl
 248 Acad Sci USA 101:7733-7738.
 249 Damen JJM, de Soet JJ, ten Cate JM. 1994. Adsorption of [3H]-lipoteichoic acid to
 250 hydroxyapatite and its effect on crystal growth. Arch Oral Biol 39(9):753-757.
 251 Eder K. 1995. Gas chromatographic analysis of fatty acid methyl esters. J Chromatogr B Biomed
 252 Appl 671(1-2):113-31.
 253 Hanson RS, Curry MV, Garner JV, Halvorson HO. 1972. Mutants of *Bacillus cereus* strain T that
 254 produce thermoresistant spores lacking dipicolinic acid have low levels of calcium. Can.
 255 J. Microbiol 18:1139-1143.
 256 Herbaud ML, Guiseppi A, Denizot F, Haiech J, Kilhoffer MC. 1998. Calcium signalling in
 257 *Bacillus subtilis*. Biochim Biophys Acta 1448(2):212-26.
 258 Huang S, Chen D, Pelczar PL, Vepachedu VR, Setlow P, Li Y. 2007. Levels of Ca²⁺-dipicolinic
 259 acid in individual *Bacillus* spores determined using microfluidic raman tweezers. J
 260 Bacteriol 189(13):4681-4687.
 261 Inui M, Suda M, Kimura S, Yasuda K, Suzuki H, Toda H, Yamamoto S, Okino S, Suzuki N,
 262 Yukawa H. 2008. Expression of *Clostridium acetobutylicum* butanol synthetic genes in
 263 *Escherichia coli*. Appl Microbiol Biotechnol 77(6):1305-16.
 264 Ito M, Terahara N, Fujinami S, Krulwich TA. 2005. Properties of motility in *Bacillus subtilis*
 265 powered by the H⁺-coupled MotAB flagellar stator, Na⁺-coupled MotPS or hybrid stators

266 MotAS or MotPB. J Mol Biol 352(2):396-408.
 267 Kidd Haack S, Garchow H, Odelson DA, Forney LJ, Klug MJ. 1994. Accuracy, reproducibility,
 268 and interpretation of fatty acid methyl ester profiles of model bacterial communities.
 269 Appl Environ Microbiol 60(7):2483–2493.
 270 Kobayashi K, Ehrlich SD, Albertini A, Amati G, Andersen KK, Arnaud M, Asai K, et al. 2003.
 271 Essential *Bacillus subtilis* genes. Proc Natl Acad Sci USA 100(8):4678-83.
 272 Koburger T, Weibezahn J, Bernhardt J, Homuth G, Hecker M. 2005. Genome-wide mRNA
 273 profiling in glucose starved *Bacillus subtilis* cells. Mol Genet Genomics 274 (1):1-12.
 274 Marvasi M, Gallagher KL, Casillas Martinez L, Molina Pagana WC, Rodríguez Santiago RE,
 275 Castilloveitia Vega G, Visscher PT. 2011. Importance of B4 medium in determining
 276 organomineralization potential of bacterial environmental isolates. Geomicrobiol J 29
 277 (10):916-924
 278 Marvasi M, Visscher PT, Perito B, Mastromei G, Casillas L. 2010. Physiological requirements
 279 for carbonate precipitation during biofilm development of *Bacillus subtilis* *etfA* mutant.
 280 FEMS Microbiol Ecology 71(3):341-350.
 281 Matsuoka H, Hirooka K, Fujita Y. 2007. Organization and function of the *YsiA* regulon of
 282 *Bacillus subtilis* involved in fatty acid degradation. J Biol Chem 282(8):5180-94.
 283 Matsuura S, Shioi JI, Imae Y, Iida S. 1979. Characterization of the *Bacillus subtilis* motile
 284 system driven by an artificially created proton motive force. J Bacteriol 140(1):28-36.
 285 Nicholson WL, Setlow P. 1990. Sporulation germination and outgrowth. In: Harwood CR,
 286 Cutting SM (ed). Molecular Biology Methods for *Bacillus*. New York: 437 p.
 287 Ordal GW, Nettleton DO, Hoch JA. 1983. Genetics of *Bacillus subtilis* chemotaxis: isolation and
 288 mapping of mutations and cloning of chemotaxis genes. J Bacteriol 154(3):1088-97.

289 Orsburn BC, Melville SB, Popham DL. 2010. EtfA catalyses the formation of dipicolinic acid in
 290 *Clostridium perfringens*. Mol Microbiol 75(1):178-186.

291 Senesi S, Ghelardi E, Celandroni F, Salvetti S, Parisio E, Galizzi A. 2004. Surface-associated
 292 flagellum formation and swarming differentiation in *Bacillus subtilis* are controlled by
 293 the *ifm* locus. J Bacteriol 186(4):1158-64.

294 Stulke J, Hillen W. 2000. Regulation of carbon catabolism in *Bacillus* species. Annu Rev
 295 Microbiol 54:849-80.

296 Thomas KJ 3rd, Rice CV. 2014. Revised model of calcium and magnesium binding to the
 297 bacterial cell wall. BioMetals 27(6). 1361-1370

298 Tojo S, Satomura T, Matsuoka H, Hirooka K, Fujita Y. 2011. Catabolite Repression of the
 299 *Bacillus subtilis* FadR Regulon, Which Is Involved in Fatty Acid Catabolism. Journal of
 300 Bacteriology 193:2388-2395.

301 Tsai MH, Saier MH Jr. 1995. Phylogenetic characterization of the ubiquitous electron transfer
 302 flavoprotein families ETF-alpha and ETF-beta. Res Microbiol 146(5):397-404.

303 Visscher PT, Dupraz C, Braissant O, Gallagher KL, Glunk C, Casillas L, Reed RES. 2010.
 304 Biogeochemistry of carbon cycling in hypersaline mats: linking the present to the past
 305 through biosignatures. In: Skeckbach J (ed). Cellular Origin, Life in Extreme Habitats
 306 and Astrobiology. Berlin:Springer Verlag. *in press*.

307 Weidenhaupt M, Rossi P, Beck C, Fischer HM, Hennecke H. 1996. *Bradyrhizobium japonicum*
 308 possesses two discrete sets of electron transfer flavoprotein genes: *fixA*, *fixB* and *etfS*,
 309 *etfL*. Arch Microbiol 165(3):169-178.

310 Wolff S, Antelmann H, Albrecht D, Becher D, Bernhardt J, Bron S, Buttner K, van Dijl JM,
 311 Eymann C, Otto A, Tam Le T, Hecker M. 2007. Towards the entire proteome of the

model bacterium *Bacillus subtilis* by gel-based and gel-free approaches. J Chromatogr B
Analyst Technol Biomed Life Sci 849(1-2):129-40.

Zhang Y, Rock CO. 2008. Membrane lipid homeostasis in bacteria. Nat Rev Micro 6:222-233.

CAPTIONS

Table I. Fatty Acid Methyl Ester (FAME) analysis for *B. subtilis* 168 and its *eftA* mutant biofilms.

Fig. 1 *In vitro* CaCO₃ precipitation on *B. subtilis* 168 and *eftA* mutant biofilms during growth in B4 medium. Strains magnification 10X. Arrows show examples of calcium carbonate crystals. *B. subtilis eftA* mutant biofilm, notice the complete absence of crystals after 8 days of incubation (Barabesi et al. 2007).

Fig. 2 Comparison of the percentage of the total amount of fatty acids of *anteiso*-C_{15:0}, *iso*-C_{17:0} and *anteiso*-C_{17:0} fatty acids produced by *B. subtilis* 168 (Δ) and its *eftA* mutant (○) during growth on B4 medium. Errors bars represent the standard deviation. Arrows indicate when CaCO₃ precipitations occur.

Fig. 3 Dipicolinic acid quantification during *B. subtilis* 168 (Δ) and its *eftA* mutant (○) biofilm formation after 7 days (168 hours) of incubation on B4 medium. Error bars represent standard deviations of three replicated samples. Arrow shows when crystals formation occurs on *B. subtilis* 168 biofilm.

Fig. 4 Comparison of lateral growth and flagella formation of *B. subtilis eftA* mutant and *B. subtilis* 168. The lower section of each figure represents the cells on the top square after flagella staining (magnification 1000X). Biofilms were produced after incubation for 5 days at 37°C

335 (1.4% agar, without calcium acetate). A) *B. subtilis etfA* mutant biofilm developed lateral growth
336 and flagella. B) No lateral growth and flagella was developed by *B. subtilis* 168. C and D) *B.*
337 *subtilis etfA* mutant and 168 growth under buffered condition (TRIS 1.2% pH 7.3). No lateral
338 growth and flagella were reported for both strains.

339